

aastex emulateapj

Submitted to ApJL

Near-infrared coronagraphy of TW Hya Trilling et al.

document

Near-infrared coronagraphic imaging of the circumstellar disk around TW Hydrae

D. E. Trilling^{1,2}, D. W. Koerner¹, J. W. Barnes^{2,3}, C. Ftaclas⁴, R. H. Brown^{2,3}

¹University of Pennsylvania, David Rittenhouse Laboratory, 209 S. 33rd Street, Philadelphia, PA 19104; trilling@hep.upenn.edu ²Visiting Astronomer at the Infrared Telescope Facility which is operated by the University of Hawaii under contract to the National Aeronautics and Space Administration ³Lunar and Planetary Laboratory, University of Arizona, Tucson, AZ 85721 ⁴Department of Physics, Michigan Technological University, 1400 Townsend Drive, Houghton, MI 49931

abstract We present ground-based near-infrared (H-band) imaging of the circumstellar disk around the nearby classical T Tauri star TW Hydrae. The scattered light image shows a face-on disk with radius 4 arcseconds (corresponding to 225 AU) and a morphology which agrees with recent images from the Hubble Space Telescope and the Very Large Array. The best fit power-law for the disk's radial surface brightness profile obeys the law $r^{-3.3 \pm 0.3}$. We use our image and published continuum flux densities to derive properties of the disk with a simple model of emission from an optically thick, flat disk. The best-fit values for disk mass and inner radius are $0.03 M_{\odot}$ and 0.3 AU; best-fit values for temperature, density, and grain opacity power law exponents (q , p , and β) are 0.7, 1.3, and 0.9, respectively. These properties are similar to those of disks around classical T Tauri stars located in more distant molecular clouds. Because of TW Hydrae's nearby location and pole-on orientation, it is a uniquely favorable object for future studies of radial disk structure at the classical T Tauri stage.

